

# Package ‘EatonEtAlChIPseq’

July 2, 2024

**Title** ChIP-seq data of ORC-binding sites in Yeast excerpted from Eaton et al. 2010

**Description** ChIP-seq analysis subset from ``Conserved nucleosome positioning defines replication origins" (PMID 20351051)

**Version** 0.42.0

**Author** Patrick Aboyoun <paboyoun@fhcrc.org>

**Maintainer** Patrick Aboyoun <paboyoun@fhcrc.org>

**Depends** GenomicRanges (>= 1.5.42), ShortRead, rtracklayer

**License** Artistic 2.0

**biocViews** ExperimentData, Saccharomyces\_cerevisiae\_Data, SequencingData, ChIPSeqData, GEO

**git\_url** <https://git.bioconductor.org/packages/EatonEtAlChIPseq>

**git\_branch** RELEASE\_3\_19

**git\_last\_commit** c17ff6c

**git\_last\_commit\_date** 2024-04-30

**Repository** Bioconductor 3.19

**Date/Publication** 2024-07-02

## Contents

orcAligns . . . . .	2
orcPeaks . . . . .	3
<b>Index</b>	<b>4</b>

---

orcAligns

*Alignments of ChIP-seq data to yeast chromosome XIV*

---

### Description

MAQ alignments to yeast chromosome XIV of ChIP-seq data of ORC-binding sites in yeast from Eaton et al. 2010

### Usage

```
data(orcAlignsRep1)
data(orcAlignsRep2)
```

### Details

This is the subset of alignments from two ChIP-seq replicates of origin recognition complex (ORC) binding to chromosome XIV of *Saccharomyces cerevisiae*. The alignments were created using MAQ (Li et al. 2008) alignment software with a maximum mismatch of 3 bases and a minimum Phred quality score of 35.

### Source

MAQ alignments extracted from [ftp://ftp.ncbi.nih.gov/pub/geo/DATA/supplementary/samples/GSM424nnn/GSM424494/GSM424494\\_wt\\_G2\\_orc\\_chip\\_rep1.mapview.txt.gz](ftp://ftp.ncbi.nih.gov/pub/geo/DATA/supplementary/samples/GSM424nnn/GSM424494/GSM424494_wt_G2_orc_chip_rep1.mapview.txt.gz) and [ftp://ftp.ncbi.nih.gov/pub/geo/DATA/supplementary/samples/GSM424nnn/GSM424494/GSM424494\\_wt\\_G2\\_orc\\_chip\\_rep2.mapview.txt.gz](ftp://ftp.ncbi.nih.gov/pub/geo/DATA/supplementary/samples/GSM424nnn/GSM424494/GSM424494_wt_G2_orc_chip_rep2.mapview.txt.gz)

### References

Conserved nucleosome positioning defines replication origins. Eaton ML, Galani K, Kang S, Bell SP, MacAlpine DM. *Genes Dev.* 2010 Apr 15;24(8):748-53.

### Examples

```
data(orcAlignsRep1)
data(orcAlignsRep2)
```

```
orcAlignsRep1
orcAlignsRep2
```

---

`orcPeaks`*Peaks from ChIP-seq alignments to yeast chromosome XIV*

---

**Description**

Peaks on yeast chromosome XIV of ChIP-seq data of ORC-binding sites in yeast from Eaton et al. 2010

**Usage**

```
data(orcPeaksRep1)
data(orcPeaksRep2)
```

**Details**

This is the subset of *Saccharomyces cerevisiae* chromosome XIV peaks from two ChIP-seq replicates of a origin recognition complex (ORC) binding experiment.

**Source**

ChIP-seq peaks extracted from [ftp://ftp.ncbi.nih.gov/pub/geo/DATA/supplementary/samples/GSM424nnn/GSM424494/GSM424494\\_wt\\_G2\\_orc\\_chip\\_rep1.bed.gz](ftp://ftp.ncbi.nih.gov/pub/geo/DATA/supplementary/samples/GSM424nnn/GSM424494/GSM424494_wt_G2_orc_chip_rep1.bed.gz) and [ftp://ftp.ncbi.nih.gov/pub/geo/DATA/supplementary/samples/GSM424nnn/GSM424494/GSM424494\\_wt\\_G2\\_orc\\_chip\\_rep2.bed.gz](ftp://ftp.ncbi.nih.gov/pub/geo/DATA/supplementary/samples/GSM424nnn/GSM424494/GSM424494_wt_G2_orc_chip_rep2.bed.gz)

**References**

Conserved nucleosome positioning defines replication origins. Eaton ML, Galani K, Kang S, Bell SP, MacAlpine DM. *Genes Dev.* 2010 Apr 15;24(8):748-53.

**Examples**

```
data(orcPeaksRep1)
data(orcPeaksRep2)
```

# Index

## \* datasets

orcAligns, [2](#)

orcPeaks, [3](#)

orcAligns, [2](#)

orcAlignsRep1 (orcAligns), [2](#)

orcAlignsRep2 (orcAligns), [2](#)

orcPeaks, [3](#)

orcPeaksRep1 (orcPeaks), [3](#)

orcPeaksRep2 (orcPeaks), [3](#)